



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

May 8, 2020

Katelynn E. King  
Regulatory Analyst  
The Scotts Company  
14111 Scottslawn Road  
Marysville, OH 43041

Subject: Registration Review Label Mitigation for Prodiamine  
Product Name: Scotts Turf Fertilizer 0.28% with Halts II  
EPA Registration Number: 538-316  
Application Dates: 6/27/19  
Decision Numbers: 553644

Dear Ms. King:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Prodiamine Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A copy of your label stamped "Accepted" is enclosed. Products shipped after 12 months from the date of this amendment must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

Page 2 of 2  
EPA Reg. No. 538-316  
Decision No. 553644

If you have any questions about this letter, please contact Darius Stanton by phone at 703-347-0433, or via email at [Stanton.darius@epa.gov](mailto:Stanton.darius@epa.gov).

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington".

Linda Arrington, Branch Chief  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

Enclosure

NOTE:

EPA Reg. No. 538-316

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

**PRODIAMINE GROUP 3 HERBICIDE**

[For preemergence control of grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns.]

[Landscape ornamentals, established perennials and wildflower plantings.]

# SCOTTS TURF FERTILIZER 0.28% WITH HALTS II

Alternate Brand Names:

[Scotts Fertilizer 22-0-8 with Halts II]

[Scotts Fertilizer 14-0-5 with Halts II]

[Scotts Fertilizer 14-2-5 with Halts II]

[Scotts Fertilizer 24-0-8 with Halts II]

[Scotts Fertilizer 0-0-7 with Halts II]

ACTIVE INGREDIENT:

Prodiamine: [N<sup>3</sup>,N<sup>3</sup>-Di-n-propyl-2, 4-dinitro-6-(trifluoromethyl)-m-phenylenediamine] ..... 0.28%

OTHER INGREDIENTS ..... 99.72%


TOTAL..... 100.00%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back/side/top/bottom [panel/label] for additional precautionary statements.

**NET WEIGHT: 50 POUNDS (22.7 kg.)**



***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye irritation. Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eyes. Call a poison control center for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or physician. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1(877)465-5161 for emergency medical treatment information.

### ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment wash water.

#### Non-Target Organism Advisory Statement:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### WEED RESISTANCE MANAGEMENT

Any weed population may contain or develop plants naturally resistant to SCOTTS TURF FERTILIZER 0.28% WITH HALTS II and other Group 3 herbicides.

Suspected herbicide-resistance weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;

NOTE:

EPA Reg. No. 538-316

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

For further information or to report suspected resistance, contact The Scotts Company at 1-800-543-8873 or at [www.scotts.com](http://www.scotts.com).

### **APPLICATION PRECAUTIONS**

Not for use on turf or plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on turf or plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

NOTE: Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

### **PRODUCT INFORMATION**

This product is a turf fertilizer and a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens), and lawns, landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development.

### **ESTABLISHED TURF**

**Application Directions:** Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation prior to weed seed germination and within 14 days of application.

**Use Precautions:** The following precautions apply to the use of this product in turf grasses and lawns.

- 1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
- 2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).
- 3) Do not cut (harvest) treated sod before 120 days after application. Do not apply newly set sod until the following year.
- 4) Application to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- 5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- 6) Do not apply to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

**Rates of Application:** This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential application may be made as long as the total amount of product applied does not exceed the maximum annual application rates listed for each turf species. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of

NOTE:

EPA Reg. No. 538-316

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

weed control.

**Maximum Annual Rates:** This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not apply more than 535 lbs. of this product per calendar year. Use higher rates of product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum yearly rate listed.

<b>MAXIMUM APPLICATION RATE/CALENDAR YEAR (by Turf Grass Species)</b>		
<b>Turf Species</b>	<b>Lbs. Product/Acre</b>	<b>Lbs. a.i./Acre</b>
Creeping Bentgrass	233	0.65
Creeping Red Fescue	267	0.75
Buffalograss Kentucky Bluegrass Perennial Ryegrass	357	1.00
Bahiagrass <sup>1</sup> Bermudagrass <sup>2</sup> Centipedegrass <sup>1</sup> Seashore Paspalum <sup>1</sup> St. Augustinegrass Tall Fescue (including turf-type) Zoysiagrass	535	1.50

<sup>1</sup>These are the maximum rates per calendar year

<sup>2</sup>May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 180 lbs/acre. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

**When to Apply after Overseeding Turf:** Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

**When to Overseed after Application:** This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results.

<b>Product Rate Lbs./Acre</b>		<b>Months Before Overwintering</b>		
<b>Lbs. Product/Acre</b>	<b>Lbs. a.i./Acre</b>	<b>North</b>	<b>Transition</b>	<b>South</b>
180	0.50	4	4	4
233	0.65	5	4	4
267	0.75	6	5	5
285	0.80	-	6	6
357	1.00	-	7	7
407	1.14	-	-	9
465	1.30	-	-	10
535	1.50	-	-	12

**Application Directions:** Apply uniformly with suitable, calibrated application equipment. Calibrate your spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. Apply this product uniformly over the area to be treated. Avoid streaking, skips and overlaps during

NOTE:

EPA Reg. No. 538-316

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

application. Calibration of spreader may be needed if conditions of area to be treated changes and rate or speed of application are changed.

<b>SPREADER RECOMMENDATIONS</b>				
<b>Spreader Settings for 1,000 sq. ft.</b>	<b>180 lbs. product/A</b>	<b>267 lbs. product/A</b>	<b>357 lbs. product/A</b>	<b>535 lbs. product/A</b>
Spreader A	(Spreader settings available at printing)			
Spreader B				
Spreader C				
(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)				

### **LANDSCAPE ORNAMENTALS**

**(Including established perennials and wildflower plantings)**

This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 5.35 to 12.28 pounds/1,000 square feet (233 to 535 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 12.28 pounds/1,000 square feet (535 pounds/acre) per year.

**Application Timing and Information:** This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

**Tolerant Ornamental Species:** This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

#### **SCIENTIFIC NAME**

*Abies spp.*  
*Abelia grandiflora*  
*Acer palmatum*  
*A. Platanoides*  
*Achillea spp.*  
*Actindia chinensis*  
*Agapanthus africanus*  
*Agapanthus orientalis*  
*Akebiaq quintata*  
*Allium cemuum*  
*Anemone hybrida*

#### **COMMON NAME**

Fir species\*\* (Balsam, Frasier, Noble, etc.)  
 Abelia: Sherwood  
 Japanese Maple  
 Norway Maple  
 Yarrow: King Edward  
 Kiwi\*  
 Lily-of-the-Nile (African Lily)  
 Five-leaf or Chocolate Vine  
 Lady's Leek, Nodding Onion  
 Japanese Anemone

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

<i>Aquilegia</i> spp.	Aquilegia: Red and Gold
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Artemesia</i> spp.	Wormwood; Silver Mound, Castle
<i>Aster</i> spp.	Aster; Bonnie Blue, Purple Dome
<i>Aster X Frikartii</i>	
<i>Athrium Filiz-femina</i>	Lady fern; Fern Lady
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Begonia</i> spp.	Fibrous Begonia: Hardy Grandis
<i>Berberis gladwynensis</i>	Barberry
<i>B. julianae</i>	Wintergreen Barberry
<i>B. mentorensis</i>	Mentor Barberry
<i>B. Thunbergii</i>	Japanese Barberry
<i>B. Verruculosa</i>	Warty Barberry
<i>Bergenia cordifolia</i>	
<i>Boltonia asteroids</i>	Snowbank
<i>Buddleia davidii</i>	Butterfly-Bush (Dward Blue); Royal Red
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Campanula carpatica</i>	Tussock Bellflower, (White Clips)
<i>Campis X Tagliabuana</i>	Trumpet Flower, Madame Galen
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Cassia artemisoides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Ceratostigma plumbagonoides</i>	
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Cleyera japonica</i>	Cleyera
<i>Citrus</i> spp.	Citrus species*
<i>Coreopsis</i> spp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
<i>Cornus florida</i>	Flowering Dogwood
<i>C. stolonifera</i>	American Dogwood
<i>Cortaderia seloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>C. buxifolius</i>	Cotoneaster
<i>C. dammeri</i>	Bearberry Cotoneaster
<i>C. microphyllus</i>	Rockspray Cotoneaster
<i>Cretaegus</i> spp.	Hawthorne
<i>Crococsmia</i> spp.	Lucifer
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Delosperma</i> spp.	Cooperi Pink
<i>Delphinium</i> spp.	Larkspur: Blue Elf
<i>Dianthus deltoides</i>	Dianthus: Maiden Pinks "Zing"
<i>D. gratianopolitanus</i>	Cheddar Pink
<i>Dodonea viscosa</i>	Hop Bush
<i>Echinacea purpurea</i>	Coneflower, Purple; Magnus
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus fortunei</i>	Wintercreeper
<i>E. japonica</i>	Evergreen Euonymus
<i>E. kiautschovica</i>	Spreading Euonymus
<i>Fatsia japonica</i>	Japanese Aralia
<i>Forsythia intermedia</i>	Border Forsythia
<i>Forsythia suspensa</i>	Weeping Forsythia
<i>F. viridissima</i>	Greenstem Forsythia

**Bold, italicized text is information for the reader and is not part of the label.**

[Bracketed information is optional text]. Text separated by / denotes and/or options.

<i>Gaillardia</i> spp.	Gaillardia, Blanket Flower: "Goblin"
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine
<i>Gaura</i> spp.	
<i>Gentiana dahurica</i>	Gentian
<i>Geranium cinereum</i>	Cranesbill
<i>Gladiolus</i> spp.	Gladiolus species**
<i>Gypsophilia repens</i>	Baby's Breath
<i>Hedera helix</i>	English Ivy
<i>Hellanthenum</i> spp.	Sunrose
<i>Hemerocallis</i> spp.	Daylily: Aztec Gold, Stella de Oro, Tender Love
<i>Hibiscus</i>	Rose of Sharon**
<i>Hibiscus</i> spp.	Mallow: Disco Belle White
<i>Hibiscus rosa-sinensis</i>	Chinese Hibiscus
<i>Hosta plantaginea</i>	Hosta, Plantain Lily (Fragrant)
<i>H. sieboldiana</i>	Hosta, "Seersucker"
<i>Houttuynia cordata</i> var. <i>variegata</i>	
<i>Hydrangea macrophylla</i>	Bigleaf Hydrangea
<i>Ilex bernyi</i>	Holly
<i>Ilex cornuta</i>	Chinese Holly
<i>I. crenata</i>	Japanese Holly
<i>I. opaca</i>	American Holly
<i>I. vomitoria</i>	Yaupon Holly, Schillings
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong
<i>Iris siberica</i>	Siberian Iris; Cabernet
<i>Iris</i> spp.	Iris species**
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact, Parsonnii
<i>J. conferta</i>	Shore Juniper; Blue Pacific
<i>J. davurica</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans</i> spp.	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica</i> X <i>Fauriei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavender</i> spp.	Lavender; Munstead
<i>Leonopodium alpinum</i>	Edelweiss
<i>Ligustrum amurense</i>	Amur. Privet
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lilium</i> spp.	Lily; Jazz
<i>Liriope muscari</i>	Big Blue Lily, Silver Mound, Evergreen Giant
<i>Liriope muscari</i> var. <i>variegata</i>	Liriope, Variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Lorapetalum chinense</i>	Burgundy
<i>Lythrum</i> spp.	Loosestrife; Modern Pink
<i>Magnolia</i> spp.	Magnolia species**
<i>Maleophora luteola</i>	Ice Plant
<i>Malus</i> spp.	Crabapple*
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

<i>Narcissus spp.</i>	Narcissus species**
<i>Nerium oleander</i>	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osmanthus heterophyllus</i>	Osmanthus (False Holly); Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fountain Grass (Drawf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado*
<i>Photinia Fraseri</i>	Frasier's Photinia (Redtip)
<i>Physotegia Virginiana</i>	Dragonhead, False; Vivid
<i>Picea spp.</i>	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-Valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. Elliottii</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine
<i>P. palustris</i>	Longleaf Pine
<i>P. radiata</i>	Monterey Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>P. taeda</i>	Loblolly Pine
<i>P. Thunbergiana</i>	Japanese Black Pine
<i>P. Virginiana</i>	Virginia Pine
<i>Pistachio spp.</i>	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>P. tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus spp.</i>	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>P. fortuneana</i>	Firethorn
<i>P. koidzumii</i>	Firethorn
<i>Pyrus spp.</i>	Bradford Pear spp.
<i>Quercus Shumardii</i>	Oak, Shumard's Red
<i>Quercus spp.</i>	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Raphiolepis umbellata</i>	Yedda Hawthorne
<i>Rhododendron</i> (including <i>Azalea</i> )	'Coral Bells', 'Formosa', 'Hino-crimson', 'PJM', 'Delaware Valley White', 'Roseum Elegans', 'Flame Creeper', 'Girard Crimson', 'Wakeilebisu', 'White Gumbo', 'George L. Tabor'
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Rudbeckia spp.</i>	Black-Eyed Susan; Goldstrum
<i>Santolina virens</i>	Santolina
<i>Saxifraga spp.</i>	Saxifrage; Purple Dome
<i>Scabiosa spp.</i>	Pincushion Flower
<i>Sedum album</i>	Stonecrop
<i>Sedum cauticola</i>	Stonecrop; Lidakense

NOTE:

EPA Reg. No. 538-316

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

<i>Sedum dasyphyllum</i>	Stonecrop
<i>Sedum spurium</i>	Stonecrop; Dragon's Blood
<i>Spiraea bumalda</i>	Spirea; Anthony Waterer
<i>Syzygium paniculatum</i>	Australian Brushcherry
<i>Taxus cuspidata</i>	Japanese Yew
<i>T. media</i>	Yew
<i>Teucrium spp.</i>	Germander
<i>Thalictrum dipterocarpum</i>	Meadow Rue
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa spp.</i>	Tulip species
<i>Veronica spp.</i>	Veronica, Speedwell; Sunny Border
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum
<i>V. plicatum</i>	Japanese Snowball
<i>V. rigidum</i>	Canary Island Viburnum
<i>V. suspensum</i>	Arrowwood Viburnum
<i>V. tinus</i>	Laurustinus
<i>V. trilobium</i>	Cranberry Bush
<i>V. wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Greater Periwinkle, Vinca
<i>Vinca minor</i>	Common Periwinkle, Vinca
<i>Vitis spp.</i>	Grape*
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet, Yucca
<i>Y. filametosa</i>	Adam's Needle
Yucca	

\* Do not use on food producing trees, vines or plants.

\*\* Not for use on container grown plants

NOTE:

EPA Reg. No. 538-316

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

## WEEDS CONTROLLED

When used in accordance with label directions, this product will provide control of the following weeds:


Barnyardgrass	Henbit	Purslane, Common
Bluegrass, Annual ( <i>Poa annua</i> )	Itchgrass	Pusley, Florida
Carpetweed	Johnsongrass (from seed)	Rescuegrass <sup>3</sup>
Chickweed, Common	Junglerice	Shepard's Purse <sup>2</sup>
Chickweed, Mouseear (from seed)		
Crabgrass (large, smooth)	Knotweed	Signalgrass, Broadleaf
Crowfootgrass	Kochia	Speedwell, Persian
Cupgrass, Woolly	Lambsquarter, Common	Sprangletop
Foxtails, Annual	Lovegrasses	Spurge, Prostrate
Goosegrass <sup>1</sup>	Panicum (Texas, fall, browntop)	Witchgrass
	Pigweed	Woodsorrel, Yellow (from seed)

<sup>1</sup> In many areas, a single application of 233 to 535 pounds/acre will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 233 to 357 pounds/acre, followed after 60 to 90 days by a second Turfgrass Species" table in the Rates of Application section of this label).

<sup>2</sup> For this weed, apply in late summer, fall or winter prior to germination.

<sup>3</sup> Suppression only.

## GERMINATION DATES



NOTE:

EPA Reg. No. 538-316

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

**FERTILIZER GUARANTEE INFORMATION**

***The following table may be used in whole or in part. Fertilizer information denoted by "X" will be added to FPL. NPK will not exceed the endpoints tested for acute toxicology studies.***

**GUARANTEED ANALYSIS**

[Product Name] [Scotts Turf Fertilizer 0.28% with Halts II]

XX-XX-XX (*insert N-P-K ratio*)

FXXX (*Insert FL Fertilizer License No.*)

Total Nitrogen .....	XX%
Soluble Potash .....	XX%
Sulfur .....	XX%
XXXXXX .....	XX%

Derived from: **insert ingredients as required to comply with State fertilizer regulations**

\*Contains X.X% slowly available ***insert source of slowly available nitrogen consistent with State fertilizer regulations***

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

***[Following are optional referral statements for Florida BMP information sources]***

**[FOR FLORIDA:**

**We recommend that you follow the Green Industries BMP's at:**

[http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/BMP\\_Book\\_final.pdf](http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/BMP_Book_final.pdf)

**[FOR FLORIDA:**

**We recommend that you follow the Golf Course BMP's at:**

<http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>

NOTE:

EPA Reg. No. 538-316

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE AND CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. **If spilled:** Sweep up and use as per label instructions or dispose of as indicated above.

NOTICE TO BUYER - Purchase of this material does not confer any rights under patents of countries outside of the United States.

## **LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as eater conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of The Scotts Company or Seller. These risks can cause: ineffectiveness of the product or other unintended consequences. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

The Scotts Company warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, The SCOTTS COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THE SCOTTS COMPANY OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF THE SCOTTS COMPANY OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHER LEGAL THEORY) RESULTING FROM THE STORAGE, USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE

NOTE:

EPA Reg. No. 538-316

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional text]. Text separated by / denotes and/or options.

OF THE PRODUCT, OR AT THE ELECTION OF THE SCOTTS COMPANY OR SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, The Scotts Company or its distributor must have prompt notice of any claim so that an immediate investigation of buyer's or user's claim can be made. Buyer and all users shall promptly notify The Scotts Company or a Scott's Company distributor or any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

EPA Reg. No. 538-316

EPA Est. No. 538-OH-1<sup>S</sup>

[Superscript used is first letter of run code on end of container]

[Manufactured by] [Sold by] [Sold and Guaranteed by]

The Scotts Company

14111 Scottslawn Rd.

Marysville, OH 43041

© 2010. Licensed by The Scotts Company LLC.

World rights reserved

[Made in USA] [Product of USA]

[Product of Country Name] [Made in Country Name]